



William Clay Albrecht, MD · Charles J. Aramburo, MD · Joey M. Bluhm, MD · Kelly S. Dempsey, MD · Elogy Etter, MD · Buckminster J. Farrow, MD · John D. Fisher, MD · Adrienne Floyd, MD · Amanda Garza · Robert M. Hillery, MD · Eddie Lambert, MD · Anne Le, MD · Liz Lee, MD · Thomas Wylie Moore, MD · Rick Ngo, MD · Guillermo Ponce de Leon, MD · Arlene Ricardo, MD · Murtaza Shakir, MD · Robert Shirley, MD · Lucrecia Sta Ana, MD · Michael Todd Wood, MD

Post-Operative instructions for Laparoscopic and Open Appendectomy

Call and schedule your post-op visit to occur approximately 7-10 days after surgery.

Wound care:

- 1) Remove outer dressing (if you have them) at 24-48 hours after surgery
- 2) Leave the steri-strips in place if you have them. They will fall off on their own.
- 3) You may have skin glue sealing your incision; this resembles super glue and is used to reinforce your dissolvable sutures making your incision waterproof and allowing you to take a shower. Do not soak in a tub or go swimming for 14 days.
- 4) You may use an ice bag to reduce swelling at the surgery site for the first 24 hours and then switch to heat if needed. It is normal to feel a knot under your surgical incision for several months after surgery.
- 5) You are able to take a shower 24 hours after surgery. Dry off the surgical incision after showering and cover with dry gauze if needed.
- 6) Antibiotics need to be taken as prescribed until they are completed.

Diet:

You may advance your diet as tolerated after surgery. If you experience nausea and vomiting after surgery start on a clear liquid diet and call the office.

Activity:

For 6 weeks after surgery

- 1) No heavy lifting over 20 lbs.
- 2) No strenuous exercise such as push-ups, sit-ups or abdominal straining
- 3) Walking is encouraged.
- 4) Normal daily activities are encouraged.
- 5) Running and biking are allowed when you feel up to it.
- 6) No driving until you are pain free and no longer using your pain medication.
- 7) Avoid extended bed rest.

Constipation and bloating:

- 1) You should have a bowel movement within 3 days of surgery. If you are constipated, increase your water intake and use an over-the-counter stool softener such as mineral oil or a laxative such as Milk of Magnesia as directed on the bottle. Pain medication often causes constipation.
- 2) If you feel bloated increase your activity.

Call our office if:

- 1) You develop any unusual signs or symptoms such as tenderness, bleeding, redness, or drainage at the surgery site.
- 2) You have a temperature of 100.5 degrees or higher.
- 3) You have shortness of breath upon exertion.

Memorial Hermann Sugar Land Location:	17510 W. Grand Parkway S., Suite 490 · Sugar Land, Texas 77479
Methodist Sugar Land Location:	16651 Southwest Freeway, Suite 360 · Sugar Land, Texas 77479
Memorial Hermann Southwest Location:	7777 Southwest Freeway, Suite 810 · Houston, Texas 77074
Memorial Hermann Katy Location:	23920 Katy Freeway, Suite 410 · Katy, Texas 77494
Memorial Hermann Memorial City Location:	920 Frostwood, Suite 620 · Houston, Texas 77024
Memorial Hermann Cypress:	27700 Northwest Freeway, Suite 330 · Cypress, Texas 77433
St. Joseph Hospital	1315 St. Joseph Pkwy Suite 1708 · Houston, TX 77002
Pearland Kirby Medical Plaza Location:	10970 Shadow Creek Parkway, Suite 220 · Pearland, Texas 77584
Woman's Hospital:	7400 Fannin Dr., Suite 850 · Houston, Texas 77054
Central Business Office:	13811 Murphy Road · Stafford, Texas 77477



William Clay Albrecht, MD · Charles J. Aramburo, MD · Joey M. Bluhm, MD · Kelly S. Dempsey, MD · Elegy Etter, MD · Buckminster J. Farrow, MD · John D. Fisher, MD · Adrienne Floyd, MD · Amanda Garza · Robert M. Hillery, MD · Eddie Lambert, MD · Anne Le, MD · Liz Lee, MD · Thomas Wylie Moore, MD · Rick Ngo, MD · Guillermo Ponce de Leon, MD · Arlene Ricardo, MD · Murtaza Shakir, MD · Robert Shirley, MD · Lucrecia Sta Ana, MD · Michael Todd Wood, MD

- 4) You are unable to urinate within 8 hours of surgery.
- 5) Your pain is not controlled with your pain medication.

Drains:

Drains are sometimes used after surgery to promote healing, you will be instructed on how to empty and document 24 hour totals on each drain (usually one and occasionally two drains are placed). This Drain Total Log should be brought to your post-op visit for your physician to review and is often one of the deciding factors regarding time for removal.

Smoking:

Smoking can cause a multitude of problems for a surgical patient, including the non-healing of a surgical incision and an increase rate of post-operative infection. Please do not smoke for two weeks prior to and two weeks after surgery.

Medication refills:

Please call your pharmacy directly and request any needed refills. The pharmacy will contact our office and refills will be expedited.

Memorial Hermann Sugar Land Location:	17510 W. Grand Parkway S., Suite 490 · Sugar Land, Texas 77479
Methodist Sugar Land Location:	16651 Southwest Freeway, Suite 360 · Sugar Land, Texas 77479
Memorial Hermann Southwest Location:	7777 Southwest Freeway, Suite 810 · Houston, Texas 77074
Memorial Hermann Katy Location:	23920 Katy Freeway, Suite 410 · Katy, Texas 77494
Memorial Hermann Memorial City Location:	920 Frostwood, Suite 620 · Houston, Texas 77024
Memorial Hermann Cypress:	27700 Northwest Freeway, Suite 330 · Cypress, Texas 77433
St. Joseph Hospital	1315 St. Joseph Pkwy Suite 1708 · Houston, TX 77002
Pearland Kirby Medical Plaza Location:	10970 Shadow Creek Parkway, Suite 220 · Pearland, Texas 77584
Woman's Hospital:	7400 Fannin Dr., Suite 850 · Houston, Texas 77054
Central Business Office:	13811 Murphy Road · Stafford, Texas 77477