



John Abdelsayed, MD · William Clay Albrecht, MD · Charles J. Aramburo, MD · Joey M. Bluhm, MD · Kelly S. Dempsey, MD · Elegy Etter, MD · Buckminster J. Farrow, MD · John D. Fisher, MD · Adrienne Floyd, MD · Amanda Garza, MD · Robert M. Hillery, MD · Eddie Lambert, MD · Anne Le, MD · Liz Lee, MD · Rick Ngo, MD · Guillermo Ponce de Leon, MD · Arlene Ricardo, MD · Robert Shirley, MD · Lucrecia Sta Ana, MD · Michael Todd Wood, MD

### **Post-Operative instructions for Hemorrhoidectomy**

**Call and schedule your post-op visit to occur approximately 7-10 days after surgery.**

#### **Wound care:**

- 1) Sitz baths are encouraged several times a day.
- 2) Moistened wipes should be used for gentle cleaning of your rectal area after a bowel movement.
- 3) You may take a bath 24 hours after surgery.
- 4) Do not sit on a donut device as this may increase swelling to the area.
- 5) Pain medication should be used as directed.
- 6) Dressings should be changed as they become wet or soiled. Gauze or sanitary napkins placed in the underwear work well.
- 7) You should start taking a stool softener daily on your first post-operative day and drink plenty of fluids. This will aid you in have a comfortable first bowel movement after surgery.

#### **Diet:**

You may advance your diet as tolerated after surgery.

#### **Activity:**

- 1) Normal daily activities are encouraged.
- 2) No driving until you are pain free and no longer using your pain medication.

#### **Constipation:**

You should have a bowel movement within 3 days of surgery. If you are constipated, increase your water intake and use an over-the-counter stool softener such as mineral oil, colace, dulcolax or a laxative such as Milk of Magnesia as directed on the bottle. Pain medication often causes constipation.

#### **Smoking:**

Smoking can cause a multitude of problems for a surgical patient, including the non-healing of a surgical incision and an increased rate of post-operative infection. Please do not smoke for two weeks prior to and two weeks after surgery.

#### **Call our office if:**

- 1) You have a temperature of 100.5 degrees or higher.
- 2) You have shortness of breath upon exertion.
- 3) You are unable to urinate within 8 hours of surgery.
- 4) Your pain is not controlled with your pain medication.
- 5) Minimal bleeding is expected. However you should report excessive bleeding.

Memorial Hermann Sugar Land Location:	17510 W. Grand Parkway S., Suite 490 · Sugar Land, Texas 77479
Methodist Sugar Land Location:	16651 Southwest Freeway, Suite 360 · Sugar Land, Texas 77479
Memorial Hermann Southwest Location:	7777 Southwest Freeway, Suite 810 · Houston, Texas 77074
Memorial Hermann Katy Location:	23920 Katy Freeway, Suite 410 · Katy, Texas 77494
Memorial Hermann Memorial City Location:	920 Frostwood, Suite 620 · Houston, Texas 77024
Memorial Hermann Cypress:	27700 Northwest Freeway, Suite 330 · Cypress, Texas 77433
St. Joseph Hospital	1315 St. Joseph Pkwy Suite 1708 · Houston, TX 77002
Pearland Kirby Medical Plaza Location:	10970 Shadow Creek Parkway, Suite 220 · Pearland, Texas 77584
Central Business Office:	13811 Murphy Road · Stafford, Texas 77477